


Amendments to the Claims:

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

Claim 1. (previously amended) A compound of formula (I)(B):



wherein

X₁ is CR₁, wherein R₁ is H, halo, amino, or nitro; and X₂ is NR₃;
R₃ is H, -SO₂ (C₁₋₆ alkyl), -SO₂ phenyl, (C=O)(C₁₋₆ alkyl), or -W'Z';
W' is a covalent bond, (C=O), SO₂, or C₁₋₆ alkyl;
Z' is C₁₋₆ alkyl, C₁₋₆ alkoxy, C₃₋₈ cycloalkyl, or a heterocyclic radical selected from the group consisting of thiazoyl, furyl, pyranyl, isobenzofuranyl, pyrrolyl, imidazolyl, pyrazolyl, isothiazolyl, isoxazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indolizinyl, isoindolyl, indolyl, indazolyl, purinyl, quinolyl, furazanyl, pyrrolidinyl, pyrrolinyl, imdazolidinyl, imidazolinyl, pyrazolidinyl, pyrazolinyl, piperidyl, piperazinyl, indolinyl, and morpholinyl; or Z' is NR₁₃R₁₄ where each of R₁₃ and R₁₄ is independently selected from C₁₋₆ alkyl, C₂₋₆ alkenyl, phenyl, benzyl, C₃₋₈ cycloalkyl, and a heterocyclic radical selected from the group consisting of thiazoyl, furyl, pyranyl, isobenzofuranyl, pyrrolyl, imidazolyl, pyrazolyl, isothiazolyl, isoxazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indolizinyl, isoindolyl, indolyl, indazolyl, purinyl, quinolyl, furazanyl, pyrrolidinyl, pyrrolinyl, imdazolidinyl, imidazolinyl, pyrazolidinyl,

pyrazolinyl, piperidyl, piperazinyl, indolinyl, and morpholinyl each of R₅, R₆, R₇ and R₈ is independently H, C ₁₋₆ alkyl, C₁₋₆ alkoxy, halo, nitro, or amino; one of R_a, R_b, R_c, R_d, and R_e is WZ and the others are independently selected from H, C ₁₋₆ alkyl, C₁₋₆ alkoxy, halo, nitro, and amino; W is -O-, O-R₉, NR₁₀, -(CO)(O)R₉, -O (CO)R₉, -(CO)NR₁₀, or -N(R₁₀)-CO-R₉, wherein R₉ is C ₁₋₆ alkylene, C ₂₋₆ alkynylene, C ₂₋₆ alkenylene, phenylene, or a heterocyclic bivalent radical selected from the group consisting of thiazoyl, furyl, pyranyl, isobenzofuranyl, pyrrolyl, imidazolyl, pyrazolyl, isothiazolyl, isoxazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indolizinyl, isoindolyl, indolyl, indazolyl, purinyl, quinolyl, furazanyl, pyrrolidinyl, pyrrolinyl, imdazolidinyl, imidazolinyl, pyrazolidinyl, pyrazolinyl, piperidyl, piperazinyl, indolinyl, and morpholinyl, and R₁₀ is H, C ₁₋₆ alkyl, C ₂₋₆ alkynyl, C ₂₋₆ alkenyl, phenyl, or a heterocyclic radical selected from the group consisting of thiazoyl, furyl, pyranyl, isobenzofuranyl, pyrrolyl, imidazolyl, pyrazolyl, isothiazolyl, isoxazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indolizinyl, isoindolyl, indolyl, indazolyl, purinyl, quinolyl, furazanyl, pyrrolidinyl, pyrrolinyl, imdazolidinyl, imidazolinyl, pyrazolidinyl, pyrazolinyl, piperidyl, piperazinyl, indolinyl, and morpholinyl; Z is a heterocyclic radical selected from the group consisting of thiazoyl, furyl, pyranyl, isobenzofuranyl, pyrrolyl, imidazolyl, pyrazolyl, isothiazolyl, isoxazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indolizinyl, isoindolyl, indolyl, indazolyl, purinyl, quinolyl, furazanyl, pyrrolidinyl, pyrrolinyl, imdazolidinyl, imidazolinyl, pyrazolidinyl, pyrazolinyl, piperidyl, piperazinyl, indolinyl, and morpholinyl, provided that when Z is pyrrolyl, piperidyl or morpholinyl the heterocyclic radical is attached through a ring

carbon; or Z is NR₁₁R₁₂ where each of R₁₁ and R₁₂ is independently selected from H, C₁₋₆ alkyl, phenyl, benzyl, C₃₋₈ cycloalkyl, and a heterocyclic radical selected from the group consisting of thiazoyl, furyl, pyranyl, isobenzofuranyl, pyrrolyl, imidazolyl, pyrazolyl, isothiazolyl, isoxazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indolizinyl, isoindolyl, indolyl, indazolyl, purinyl, quinolyl, furazanyl, pyrrolidinyl, pyrrolinyl, imidazolidinyl, imidazolinyl, pyrazolidinyl, pyrazolinyl, piperidyl, piperazinyl, indolinyl, and morpholinyl; or NR₁₁R₁₂ taken together is a C₆₋₈ cycloalkylimino radical; each of the above hydrocarbyl or heterocyclic groups being optionally substituted with between 1 and 3 substituents selected from C₁₋₃ alkyl, C₁₋₃ alkoxy, halo, hydroxy, phenyl, and phenyl(C₁₋₃ alkyl); and wherein each of the above heterocyclic groups may be attached to the rest of the molecule by a carbon atom or a heteroatom; provided that R_b, R_d, R₅, R₆, R₇ and R₈, if halo, are selected from chloro; or a pharmaceutically acceptable salt, amide, ester, or hydrate thereof.

2. (original) A compound of claim 1, wherein R₃ is H or C₁₋₃ alkyl.
3. (original) A compound of claim 1, wherein R₃ is -(C=O)C₁₋₆ alkyl.
4. (original) A compound of claim 1, wherein R₃ is -SO₂(C₁₋₃ alkyl).
5. (original) A compound of claim 4 wherein R₃ is methylsulfonyl.
6. (original) A compound of claim 1, wherein W' is a covalent bond.
7. (original) A compound of claim 1, wherein W' is SO₂ or (C=O).

8. (original) A compound of claim 1, wherein R_c is WZ.
9. (original) A compound of claim 1, wherein R_b or R_d is WZ.
10. (original) A compound of claim 1, wherein W is ethoxy, propoxy, or butoxy.
11. (original) A compound of claim 1, wherein W is -O-.
12. (original) A compound of claim 1, wherein one of R_b, R_c, and R_e is WZ and the others are independently selected from H, methyl, ethyl, methoxy, ethoxy, amino, nitro, and halo; and R_a and R_d are each independently H or methyl.
13. (original) A compound of claim 1, wherein at least two of the following apply: R_c is WZ; W is propoxy or ethoxy; and Z is N-piperidino, 2-(N-methyl)pyrrolidino, or N,N-dimethyl.
14. (previously amended) A compound of claim 1, wherein Z is pyrrolidino, N-methyl-pyrrolidino, pyridyl, thiazoyl, piperidino, or NR₁₁R₁₂ where each of R₁₁ and R₁₂ is independently selected from H, C₁₋₆ alkyl, phenyl, benzyl, C₃₋₆ cycloalkyl, and a heterocyclic radical selected from the group consisting of thiazoylyl, furyl, pyranyl, isobenzofuranyl, pyrrolyl, imidazolyl, pyrazolyl, isothiazolyl, isoxazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indolizinyl, isoindolyl, indolyl, indazolyl, purinyl, quinolyl, furazanyl, pyrrolidinyl, pyrrolinyl, imdazolidinyl, imidazolinyl, pyrazolidinyl, pyrazolinyl, piperidyl, piperazinyl, indolinyl, and morpholinyl or taken together with the N form a C₆₋₈ cycloalkylamino radical.

15. (previously amended) A compound of claim 1, wherein one of R_b , R_c , and R_e is WZ and the others are independently selected from H, methyl, ethyl, methoxy, ethoxy, amino, and halo; and R_a and R_d are each independently H or methyl;
W is -O- or C₁₋₃ alkoxy;
Z is pyrrolidino, N-methylpyrrolidino, pyridyl, thiazoyl, piperidino, piperazino, N-methylpiperazino, or NR₁₁R₁₂ where each of R₁₁ and R₁₂ is independently selected from H, C₁₋₂ alkyl, phenyl, benzyl, C₃₋₈ cycloalkyl, and a heterocyclic radical selected from the group consisting of thiazoylyl, furyl, pyranyl, isobenzofuranyl, pyrrolyl, imidazoyl, pyrazoyl, isothiazoyl, isoxazoyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indolizinyl, isoindolyl, indolyl, indazolyl, purinyl, quinolyl, furazanyl, pyrrolidinyl, pyrrolinyl, imidazolidinyl, imidazolinyl, pyrazolidinyl, pyrazolinyl, piperidyl, piperazinyl, indolinyl, and morpholinyl; each of R₆ and R₇ are each independently H, methyl, methoxy, or ethoxy; each of R₅ and R₈ is H.
16. (original) A compound of claim 15, wherein R₃ is H or -SO₂(C₁₋₆ alkyl).
17. (original) A compound of claim 15, wherein R₃ is SO₂(phenyl) and (C=O)(C₁₋₆ alkyl).
18. (previously amended) A compound of claim 15, selected from 2-[4-[2-[1-(methyl)-2-pyrrolidinyl]ethoxy]phenyl]-1H-indole, 2-[4-[2-[1-(methyl)-2-pyrrolidinyl]ethoxy]phenyl]-1-(methylsulfonyl)-1H-indole, 2-(4-(3-(4-methylpiperazino)propoxy)-phenyl)indole; and 1-(methylsulfonyl)-2-(4-(3-(4-methylpiperazino)-propanoxy)phenyl)indole; or a pharmaceutically acceptable salt, amide, ester, or hydrate thereof.

19. (cancelled)
20. (original) A pharmaceutical composition comprising a compound of formula (I)B and a pharmaceutically acceptable carrier.
21. (previously amended) A pharmaceutical composition of claim 20, wherein said compound has a formula wherein: one of R_b, R_c, and R_e is WZ and the others are independently selected from H, methyl, ethyl, methoxy, ethoxy, amino, and halo; R_a and R_d are each independently H or methyl; W is -O- or C₁₋₃ alkoxy; Z is pyrrolidino, N-methylpyrrolidino, pyridyl, thiazoyl, piperidino, or NR₁₁R₁₂ where each of R₁₁ and R₁₂ is independently selected from H, C₁₋₂ alkyl, phenyl, benzyl, C₃₋₈ cycloalkyl, and a heterocyclic radical selected from the group consisting of thiazoylyl, furyl, pyranyl, isobenzofuranyl, pyrrolyl, imidazolyl, pyrazolyl, isothiazolyl, isoxazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indolizinyl, isoindolyl, indolyl, indazolyl, purinyl, quinolyl, furazanyl, pyrrolidinyl, pyrrolinyl, imdazolidinyl, imidazolinyl, pyrazolidinyl, pyrazolinyl, piperidyl, piperazinyl, indolinyl, and morpholinyl; and R₆ and R₇ are each independently H, methyl, methoxy, or ethoxy.
22. (previously amended) A pharmaceutical composition of claim 21, wherein said compound has a formula selected from 2-[4-[2-[1-(methyl)-2-pyrrolidinyl]ethoxy]phenyl]-1H-indole; 2-[4-[2-[1-(methyl)-2-pyrrolidinyl]ethoxy]phenyl]-1-(methylsulfonyl)-1H-indole; 2-[3-[3-Piperidinopropoxy]phenyl]-1-(methylsulfonyl)-1H-indole; 2-(4-(3-(4-methylpiperazino)propoxy)-phenyl)indole; and

1-(methylsulfonyl)-2-(4-(3-(4-methylpiperazino)-
propoxy)phenyl)indole; or a pharmaceutically acceptable salt,
amide, ester, or hydrate thereof.

23. (cancelled)
24. (cancelled)
25. (cancelled)
26. (cancelled)
27. (cancelled)
28. (cancelled)
29. (cancelled)
30. (cancelled)
31. (cancelled)
32. (cancelled)
33. (new) A compound of claim 1 selected from 2-[4-[1-(methyl)-4-piperidinyl]oxyphenyl]-1-(methylsulfonyl)-1H-indole;
2-[4-[1-(methyl)-4-piperidinyl]oxyphenyl]-1H-indole;
2-[4-[3-dimethylaminopropoxy]phenyl]-1-(methylsulfonyl)-1H-indole;
2-[4-[3-dimethylaminopropoxy]phenyl]-1H-indole;
2-[4-[4-pyridinyl]methoxyphenyl]-1-(methylsulfonyl)-1H-indole;
2-[4-[2-diethylaminoethoxy]phenyl]-1-(methylsulfonyl)-1H-indole;

2-[4-[2-diethylaminoethoxy]phenyl]-1H-indole;
2-[4-[2-(2-pyridinyl)ethoxyphenyl]-1-(methylsulfonyl)-1H-indole;
and 2-[4-[2-(2-pyridinyl)ethoxyphenyl]-1H-indole; or a
pharmaceutically acceptable salt, amide, ester, or hydrate thereof.

34. (new) A pharmaceutical composition of claim 20, wherein said compound has a formula selected from 2-[4-[1-(methyl)-4-piperidinyl]oxyphenyl]-1-(methylsulfonyl)-1H-indole;
2-[4-[1-(methyl)-4-piperidinyl]oxyphenyl]-1H-indole;
2-[4-[3-dimethylaminopropoxy]phenyl]-1-(methylsulfonyl)-1H-indole;
2-[4-[3-dimethylaminopropoxy]phenyl]-1H-indole;
2-[4-[4-pyridinyl]methoxyphenyl]-1-(methylsulfonyl)-1H-indole;
2-[4-[2-diethylaminoethoxy]phenyl]-1-(methylsulfonyl)-1H-indole;
2-[4-[2-diethylaminoethoxy]phenyl]-1H-indole;
2-[4-[2-(2-pyridinyl)ethoxyphenyl]-1-(methylsulfonyl)-1H-indole;
and 2-[4-[2-(2-pyridinyl)ethoxyphenyl]-1H-indole; or a
pharmaceutically acceptable salt, amide, ester, or hydrate thereof.